

Letters to the Editor

The Board of Editors does not hold itself responsible for opinions expressed in the letters published in this section. The notes containing short reports of original investigations communicated to this section should not contain many figures and should not exceed 500 words in length. The contributions reaching the Secretary by the 15th of any month may be expected to appear in the issue for the next month. No proof will be sent to the author.

5

ON THE CLASSICAL RADIATION BY AN ELECTRON IN UNIFORM CIRCULAR MOTION IN THE FORM OF MULTIPOLES

T. P. KHAN

D. A. COLLEGE, GARIA, 24-PARGANAS, WEST BENGAL, INDIA.

(Received September 16, 1966).

INTRODUCTION

The problem of classical radiation by electron in uniform circular motion was extensively discussed by Schott (1912). While Schott's interest was to explain atomic structure Schwinger (1912) repeated that analysis of the same problem to find out the maximum energy limit to be obtained by synchro-cyclotron. In this paper attempt has been made to obtain classical radiation by electron in uniform circular motion in the form of multipoles. An electric charge within the medium of dielectric constant ϵ is moving in a circular orbit of radius R with constant angular velocity ω_0 . The motion is assumed to be confined in a plane and the angular distributions of the radiation have been obtained after integration of the usual energy term on the surface of a sphere.

Spherical wave solution of Maxwell's Equation :

The Maxwell's equations for a source moving in a circular orbit of radius R and constant angular velocity ω_0 in the medium of dielectric constant ϵ ,

$$\nabla^2 \phi - \frac{\epsilon}{c^2} \frac{\partial^2 \phi}{\partial t^2} = \frac{4\pi}{\epsilon} \rho \quad \dots (1)$$

$$\nabla^2 A - \frac{\epsilon}{c^2} \frac{\partial^2 A}{\partial t^2} = \frac{4\pi}{c} J \quad \dots (2)$$

where ϕ , A , ρ , J have their usual meaning. Out of these two equations (1) and

(2) solution of single equation is enough to determine the two quantities ϕ , A , since ϕ , A are related by the Gauge

$$A = \frac{\epsilon v}{c} \phi$$

The potentials ϕ , A , and the source density $\rho(x)$, $J(x)$ are transformed in space and time according to the general rule of Fourier transforms.

$$F(x, t) = \frac{1}{(2\pi)^2} \int d^3K \int d\omega F(K, \omega) e^{ik \cdot x - i\omega t}$$

and the equations (1) and (2) are transformed into

$$\left[K^2 - \frac{\epsilon \omega^2}{c^2} \right] \phi(K, \omega) = \frac{4\pi}{\epsilon} \rho(K, \omega) \quad \dots (3)$$

$$\left[K^2 - \frac{\epsilon \omega^2}{c^2} \right] A(K, \omega) = \frac{4\pi}{C} J(K, \omega) \quad \dots (4)$$

with reference to a spherical polar co-ordinate system we assign the position of the charged particle at any instant to be at $(R, \omega_0 t, \pi/2)$ while the position of observation point has the co-ordinate (r, ϕ, θ) .

From the knowledge of Fourier transforms (Morse *et al*).

$$\rho(K, \omega) = \frac{e}{2\pi} \int e^{-ikR[\sin u \cos(\omega_0 t - v)] - i\omega t} dt \quad \dots (5)$$

substituting (5) in (3) and applying the same technique of Fourier transforms, we obtain

$$\begin{aligned} \phi(\omega) &= \frac{2e}{\epsilon(\omega)} \frac{1}{(2\pi)^{3/2}} \int \int e^{iK \cdot x - iKR[\sin u \cos(\omega_0 t - v)] - i\omega t} d^3K dt \\ &= \frac{2e}{\epsilon(\omega)} \frac{1}{(2\pi)^{3/2}} \sum \sum \alpha_{mn} \delta(\omega - m\omega_0) e^{im\phi} \rho_n^m(\cos \theta) j_n \left(\frac{\sqrt{\epsilon\omega}}{c} R \right) h_n \left(\frac{\sqrt{\epsilon\omega}}{c} r \right) \dots (6) \end{aligned}$$

Multipole Expansion of the Electromagnetic fields

The components of the Electromagnetic fields in (r, ϕ, θ) directions

$$E_r = -\frac{\partial \phi}{\partial r} \quad \because A_r = 0$$

$$E_\phi = -\frac{1}{C} \frac{\partial A_\phi}{\partial t} - \frac{1}{r} \frac{\partial \Phi}{\partial \phi} = -\frac{\epsilon v}{c^2} \frac{\partial \Phi}{\partial t} - \frac{1}{r} \frac{\partial \Phi}{\partial \phi}$$

$$E_\theta = -\frac{1}{r \sin \phi} \frac{\partial \Phi}{\partial \theta} \quad \because A_\theta = 0$$

$$H_r = -\frac{1}{r \sin \phi} \frac{\partial A \phi}{\partial \theta} = -\frac{c}{C} \frac{v}{r \sin \phi} \frac{\partial \Phi}{\partial \theta}$$

$$H_\phi = 0$$

$$H_\theta = \frac{1}{r} \frac{\partial}{\partial r} (r A \phi) = \frac{ev}{cr} \frac{\partial}{\partial r} (r \Phi)$$

From (6) $\Phi(t)$ is obtained from the rule of Fourier transform

$$\Phi(t) = \frac{1}{\sqrt{2\pi}} \int \Phi(\omega) e^{-i\omega t} d\omega \quad \text{as}$$

$$\Phi(t) = \frac{2e}{c} \frac{1}{(2\pi)^2} \sum \sum \alpha_{mn} e^{im\phi} \rho_n^m(\cos \theta) e^{-im\omega_0 t} j_n(\sqrt{\epsilon m \beta}) h_n \left(\frac{\sqrt{\epsilon m \omega_0} r}{c} \right)$$

For energy transfer in the radial direction we require only E_ϕ and H_θ , which are when expanded in multipoles,

$$E_\phi n, m = +im \left[\frac{ev}{c^2} \omega_0 - \frac{1}{r} \right] \phi_{nm}(t)$$

$$\sum \sum E_{nm} = \frac{2eim}{c(2\pi)^2} \left[\frac{ev}{c^2} \omega_0 - \frac{1}{r} \right] \sum \sum \alpha_{mn} e^{-im\omega_0 t} \times e^{im\phi} P_n^m(\cos \theta)$$

$$\times j_n(\sqrt{\epsilon m \beta}) e^{i\sqrt{\epsilon m \omega_0} r / c} \frac{e^{i\sqrt{\epsilon m \omega_0} r / c}}{\sqrt{\epsilon m \omega_0}} \frac{1}{r} \quad \dots \quad (7)$$

$$H_{\theta nm} = \frac{ev}{c} \frac{\partial}{\partial r} \phi_{nm}(t) \text{ neglecting other terms.}$$

$$\sum \sum H_{nm} = \frac{ev}{c} \frac{\partial}{\partial r} \sum \sum \phi_{nm}(t) \alpha_{mn} \quad \dots \quad (8)$$

Angular distribution of multipole radiation

The electromagnetic fields as obtained from (7) and (8) when terms upto $\left(\frac{1}{r}\right)$ are retained

$$E_\phi \rightarrow \frac{2e}{\sqrt{\epsilon}} \cdot \frac{\beta}{(2\pi)^2} \sum \sum \alpha_{mn} e^{im\phi} \frac{e^{i\sqrt{\epsilon m \omega_0} r / c - im\omega_0 t}}{r} j_n(\sqrt{\epsilon m \beta}) e^{im\phi} P_n^m(\cos \theta)$$

$$H_\theta \rightarrow 2e \cdot \frac{\beta}{(2\pi)^2} \sum \sum \alpha_{mn} e^{im\phi} \frac{e^{i\sqrt{\epsilon m \omega_0} r / c - im\omega_0 t}}{r} j_n(\sqrt{\epsilon m \beta}) e^{im\phi} P_n^m(\cos \theta)$$

The time averaged power radiated per unit solid angle is

$$\frac{d\rho}{d\Omega} = \frac{e^2\beta^2}{\sqrt{\epsilon(\pi)^2}} j_n^2(\sqrt{\epsilon m}\beta) |X_{m,n}(\theta, \phi)|^2$$

The table lists some of the simpler angular distributions.

| $X_{m,n}(\theta, \phi)^2$ | | |
|---------------------------|---|-----------------------------------|
| m | | |
| | ± 1 | ± 2 |
| 1 Dipole | $\frac{3}{16\pi}(1+\cos^2\theta)$ | |
| 2 Quadrupole | $\frac{5}{16}(1-3\cos^2\theta$ $+4\cos^4\theta)$ | $\frac{5}{16\pi}(1-\cos^4\theta)$ |

The author expresses thanks to Dr. T. Roy of Jadavpur University for guidance and encouragement.

REFERENCES

- Jackson, *Classical Electrodynamics*, John Wiley & Sons, Inc., Page 550.
 Morse & Feshbach *Methods of Theoretical Physics*, : I & II, McGRAW—HILL BOOK Company Inc., Page 621, 827, 1325, 1362.
 Schwinger, 1949, *On Classical Radiation of accelerated electrons*, *Phys. Rev.* **75**, 1912.
 Schott, 1912, *Electromagnetic Radiation*, Cambridge University Press,